

Effect of the δ -meson on the instabilities of nuclear matter under strong magnetic fields

A. Rabhi,^{1,2,*} C. Providência,^{1,†} and J. Da Providência^{1,‡}

¹*Centro de Física Computacional, Department of Physics,
University of Coimbra, 3004-516 Coimbra, Portugal*

²*Laboratoire de Physique de la Matière Condensée,
Faculté des Sciences de Tunis, Campus Universitaire, Le Belvédère-1060, Tunisia*

(Dated: September 6, 2009)

We study the influence of the isovector-scalar meson on the spinodal instabilities and the distillation effect in asymmetric non-homogenous nuclear matter under strong magnetic fields, of the order of $10^{18} - 10^{19}$ G. Relativistic nuclear models both with constant couplings (NLW) and with density dependent parameters (DDRH) are considered. A strong magnetic field can have large effects on the instability regions giving rise to bands of instability and wider unstable regions. It is shown that for neutron rich matter the inclusion of the δ meson increases the size of the instability region for NLW models and decreases it for the DDRH models. The effect of the δ meson on the transition density to homogeneous β -equilibrium matter is discussed. The DDRH δ model predicts the smallest transition pressures, about half the values obtained for NL δ .

PACS numbers: 21.65.-f 26.60.Kp 26.60.-c 97.60.Jd

Neutron stars with very strong magnetic fields of the order of $10^{14} - 10^{15}$ G are known as magnetars [1–3] and they are believed to be the sources of the intense gamma and X rays detected in 1979 (for a review refer to [4]). To date, 16 magnetars have been identified as short γ -ray repeaters or anomalous X-ray pulsars, although some are still unconfirmed candidates [5]. However, according to Ref. [6], a fraction as high as 10% of the neutron star population could be magnetars. These neutron stars are hot, young stars, ~ 1 kyear old. In [7], Thompson and Duncan have considered turbulent dynamo amplifications in young neutron stars as a mechanism for generating the strong magnetic fields.

In magnetars the crust is stressed by very strong forces which deform the crust and may crack it. Once the surface cracks, the violent motions blast particles along the magnetic fields, triggering gamma rays and x-rays. In particular, scientists believe that the giant burst of energy experienced by the magnetar SGR 1806-20 in 2004 was triggered by a "starquake" in the neutron star's crust that caused a catastrophic disruption in the magnetar's magnetic field. SGR 1806-20 is the most magnetic object observed and has on the surface a magnetic field of intensity over 10^{15} G [5].

In [8] the authors have shown how to obtain both the moment of inertia of a neutron star as well as the fractional moment of inertia of its crust only in terms of the mass and the radius of the star. For the fractional moment of inertia of the crust has an additional dependence of the equation of state (EOS) which enters through the values of the pressure and the density at the crust core

transition [8, 9]. The transition density enters just as a correction to the fractional moment of inertia, while the transition pressure is the main EOS parameter which defines that quantity. Using this result, Link *et al.* [9] have obtained a lower limit for the neutron star radius with a given mass, from the glitches occurring in the Vela pulsar and in other six pulsars. This constraint will put severe restrictions on the acceptable equations of state of stellar matter if the radius and mass of a neutron star is measured.

Spin-up glitches have also been observed in all known persistent AXP [10, 11]. However, different from the conventional low-field radio pulsar, the glitches in AXPs are accompanied by a significant recovery of the spin down rate of the pulsar. It would be interesting if information of the crust and star properties could also be obtained from these glitches.

It was recently shown in Ref. [12] that a strong magnetic field, of the order of $10^{18} - 10^{19}$ G, has large effects on the instability regions of nuclear matter. Relativistic nuclear models both with constant couplings and with density dependent parameters were considered. It was shown that a strong magnetic field can have large effects on the thermodynamic spinodal instabilities zones giving rise to bands of instability and wider unstable regions. As a consequence, it was predicted larger transition densities at the inner edge of the crust of compact stars with a strong magnetic field. The direction of instability gives rise to a very strong distillation effect if protons occupy only partially a Landau level. However, for almost full Landau levels an anti-distillation effect may occur.

In this paper, which completes Ref. [12], we study the influence of the isovector scalar meson on the low density instabilities of asymmetric nuclear matter under strong magnetic fields. We estimate the density and pressure at the transition from the non-homogeneous to the homogeneous phase in stellar matter from the crossing of the

*Electronic address: rabhi@teor.fis.uc.pt

†Electronic address: cp@teor.fis.uc.pt

‡Electronic address: providencia@teor.fis.uc.pt

EOS in β -equilibrium matter with the spinodal and discuss the effect of the magnetic field on the direction of instability. We consider two kinds of relativistic mean-field approaches: non-linear Walecka models (NLW) models with constant coupling parameters and density dependent relativistic hadronic (DDRH) models with density-dependent coupling parameters. The last models seem to give more realistic results at subsaturation densities [13]. The inclusion of the δ -meson brings to the isovector channel the same symmetry existing already in the isoscalar channel with the meson pair (σ, ω) responsible for saturation in RMF models [14, 15]. The presence of the δ -meson softens the symmetry energy at subsaturation densities and hardens it above saturation density, giving rise to stable compact stars with larger masses [16]. In [13] it was shown that the instability region in the isovector direction becomes smaller if the δ meson is included. Therefore, it is expected that in the presence of a strong magnetic field these differences become larger.

In the present paper we consider the NLW models $NL\rho$ and $NL\delta$ [15] and the DDRH models TW [17] and $DDRH\delta$ [18]. Only $NL\delta$ and $DDRH\delta$ include the δ -meson.

For the description of the neutron star matter, we employ the standard mean-field theory (MFT) approach. A complete set of the equations and the description of the method can be found in Ref. [12, 19]. The Lagrangian density of TW [17, 20] and $DDRH\delta$ [18, 21] models reads:

$$\begin{aligned} \mathcal{L} = & \bar{\Psi}_b \left(i\gamma_\mu \partial^\mu - q_b \gamma_\mu A^\mu - m_b + \Gamma_\sigma \sigma + \Gamma_\delta \vec{\tau}_b \cdot \vec{\delta} \right. \\ & - \Gamma_\omega \gamma_\mu \omega^\mu - \frac{1}{2} \Gamma_\rho \tau_{3b} \gamma_\mu \rho^\mu - \frac{1}{2} \mu_N \kappa_b \sigma_{\mu\nu} F^{\mu\nu} \left. \right) \Psi_b \\ & + \frac{1}{2} \partial_\mu \sigma \partial^\mu \sigma - \frac{1}{2} m_\sigma^2 \sigma^2 + \frac{1}{2} \partial_\mu \vec{\delta} \partial^\mu \vec{\delta} - \frac{1}{2} m_\delta^2 \vec{\delta}^2 \\ & + \frac{1}{2} m_\omega^2 \omega_\mu \omega^\mu - \frac{1}{4} \Omega^{\mu\nu} \Omega_{\mu\nu} \\ & - \frac{1}{4} F^{\mu\nu} F_{\mu\nu} + \frac{1}{2} m_\rho^2 \rho_\mu \rho^\mu - \frac{1}{4} P^{\mu\nu} P_{\mu\nu}, \end{aligned} \quad (1)$$

where Ψ_b are the baryon ($b=n, p$) Dirac fields; σ , ω , ρ , and δ represent the scalar, vector, isovector-vector and isovector-scalar meson fields, which are exchanged for the description of nuclear interactions and $A^\mu = (0, 0, Bx, 0)$ refers to a constant external magnetic field along the z-axis. The nucleon mass and isospin projection for the protons and neutrons are denoted by m_b and $\tau_{3b} = \pm 1$, respectively. The mesonic and electromagnetic field strength tensors are given by their usual expressions: $\Omega_{\mu\nu} = \partial_\mu \omega_\nu - \partial_\nu \omega_\mu$, $P_{\mu\nu} = \partial_\mu \rho_\nu - \partial_\nu \rho_\mu$, and $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$. The nucleon anomalous magnetic moments (AMM) are introduced via the coupling of the baryons to the electromagnetic field tensor with $\sigma_{\mu\nu} = \frac{i}{2} [\gamma_\mu, \gamma_\nu]$ and strength κ_b with $\kappa_n = g_n/2 = -1.91315$ for the neutron and $\kappa_p = (g_p/2 - 1) = 1.79285$ for the proton, respectively. The electromagnetic field is assumed to be externally generated (and thus has no associated field equation), and only frozen-field configurations will be considered. The density dependent strong




FIG. 1: Symmetry energy for all models used in the present work.

interaction couplings are denoted by Γ , the electromagnetic couplings by q and the nucleon, mesons masses by m . The density dependent coupling parameters are adjusted in order to reproduce some of the nuclear matter bulk properties using the following parametrization

$$\Gamma_i(\rho) = \Gamma_i(\rho_{sat}) f_i(x), \quad i = \sigma, \omega, \rho, \delta \quad (2)$$

where $x = \rho/\rho_{sat}$, with

$$f_i(x) = a_i \frac{1 + b_i(x + d_i)^2}{1 + c_i(x + d_i)^2}, \quad i = \sigma, \omega, \quad (3)$$

for TW,

$$f_\rho(x) = \exp[-a_\rho(x - 1)], \quad (4)$$

and,

$$f_i(x) = a_i \exp[-b_i(x - 1)] - c_i(x - d_i), \quad i = \rho, \delta \quad (5)$$

for $DDRH\delta$, with the values of the parameters m_i , Γ_i , a_i , b_i , c_i and d_i , $i = \sigma, \omega, \rho, \delta$ given in Table I.

In the sequel, we define the magnetic field in units of the electron critical field $B_e^c = 4.414 \times 10^{13}$ G, so that $B = B^* B_e^c$.

For the $DDRH\delta$ model we use the parametrization given in [18, 21] except for the parameter $\Gamma_\rho(\rho_{sat})$ which we increase so that the symmetry energy is 31 MeV at saturation density and not 25 MeV as in [18], see Table I. The inclusion of the δ meson reduces the symmetry energy at subsaturation densities but makes it much stiffer at supra-saturation densities because the δ meson field reaches saturation and the ρ -meson field increases always with density [14, 15]. This is seen in Fig. 1 where the symmetry energy is given for the all models we are studying.

For the NLW models, *i.e.* $NL\rho$ and $NL\delta$, we add to the Lagrangian density, Eq. (1), with $g_i = \Gamma_i$, the scalar

i	m_i (MeV)	Γ_i	a_i	b_i	c_i	d_i
σ	550	10.72854	1.365469	0.226061	0.409704	0.901995
ω	783	13.29015	1.402488	0.172577	0.344293	0.983955
TW						
ρ	763	7.32196	0.515			
DDRH δ						
ρ	763	12.7530	0.095268	2.171	0.05336	17.8431
δ	980	7.58963	0.01984	3.4732	-0.0908	-9.811

TABLE I: Parameters for the TW and DDRH δ models. These two models have the same parametrization for the σ and ω mesons.

meson self-interaction terms

$$\mathcal{L}_{nl} = -\frac{1}{3}bm_n(g_\sigma\sigma)^3 - \frac{1}{4}c(g_\sigma\sigma)^4,$$

where b and c are two dimensionless parameters. The coupling parameters are constant and given in Ref. [15].

The stability conditions for asymmetric nuclear matter, keeping volume and temperature constant, are obtained from the free energy density $\mathcal{F}$, imposing that this function is a convex function of the densities ρ_p and ρ_n , *i.e.* the symmetric matrix with the elements [22, 23]

$$\mathcal{F}_{ij} = \left(\frac{\partial^2 \mathcal{F}}{\partial \rho_i \partial \rho_j} \right)_T \quad (6)$$

is positive. At zero temperature, the free energy density coincides with the energy density. We define the thermodynamic spinodal at $T = 0$ as the curve on the (ρ_n, ρ_p) plane for which the determinant of the $\mathcal{F}_{ij}$ is zero. The eigenvalues of the stability matrix are given by

$$\lambda_{\pm} = \frac{1}{2} \left(\text{Tr}(\mathcal{F}) \pm \sqrt{\text{Tr}(\mathcal{F})^2 - 4\text{Det}(\mathcal{F})} \right) \quad (7)$$

and the eigenvectors $\delta\rho_{\pm}$ by

$$\frac{\delta\rho_i^{\pm}}{\delta\rho_j^{\pm}} = \frac{\lambda_{\pm} - \mathcal{F}_{jj}}{\mathcal{F}_{ji}}, \quad i, j = p, n. \quad (8)$$

In Fig. 2 we show the spinodal sections for several magnetic field intensities and for two NLW models: NL ρ without δ -meson (thin lines) and NL δ including δ (thick lines). As discussed in [12], due to the Landau quantization, the magnetic field has a strong effect not only on the size, but also on the shape of the spinodal zone. Unlike the $B = 0$ case, the instability zone is no longer symmetric with respect to the $\rho_n = \rho_p$ line. For the proton-rich matter, the inclusion of the δ -meson has only a small effect on the spinodal, namely a small reduction similar to the effect already described for the $B = 0$ [24]. For neutron-rich matter in the presence of a strong magnetic field, the effect of δ -meson is stronger: the NL δ model has a spinodal region larger than NL ρ . This effect is similar whether the AMM is included or not.

The DDRH models behave in a different way due to density dependence of the coupling parameters. In Fig. 3,

TABLE II: Predicted density, proton fraction and pressure at the inner edge of the crust of a compact star at zero temperature, as defined by the crossing between the thermodynamic instability region of np matter and the β -equilibrium EOS for homogeneous, neutrino-free stellar matter in the (ρ_p, ρ_n) plane. The AMM is not included.

B^*	Model	$\rho_b^{\text{cross}}(\text{fm}^{-3})$	Y_p	$P_m(\text{MeVfm}^{-3})$
0	NL ρ	0.067	0.013	0.255
	NL δ	0.063	0.010	0.175
	TW	0.085	0.037	0.523
	DDRH δ	0.086	0.036	0.269
3×10^4	NL ρ	0.071	0.043	0.284
	NL δ	0.071	0.040	0.264
	TW	0.071	0.056	0.342
	DDRH δ	0.077	0.056	0.259
5×10^4	NL ρ	0.084	0.068	0.467
	NL δ	0.084	0.064	0.467
	TW	0.082	0.086	0.461
	DDRH δ	0.084	0.086	0.306
10^5	NL ρ	0.105	0.121	0.822
	NL δ	0.105	0.117	0.896
	TW	0.101	0.146	0.673
	DDRH δ	0.096	0.148	0.404
2×10^5	NL ρ	0.128	0.207	1.203
	NL δ	0.130	0.203	1.447
	TW	0.128	0.236	1.016
	DDRH δ	0.118	0.237	0.644
5×10^5	NL ρ	0.162	0.367	1.275
	NL δ	0.170	0.369	1.830
	TW	0.198	0.402	2.331
	DDRH δ	0.228	0.422	4.460

we present the results of the thermodynamic spinodal instability region for TW (thin lines) and DDRH δ (thick lines). For $B^* = 10^5$ and 2×10^5 the spinodal has three and two bands, see Fig. 3, corresponding to the occupation of the first three and two Landau levels, respectively. For proton-rich matter with $B^* = 10^5$ and 2×10^5 , the inclusion of the δ -meson reduces the size of the spinodal zone corresponding to the last occupied Landau level (LL), *i.e.*, third LL for $B^* = 10^5$ and second LL for $B^* = 2 \times 10^5$. The other instabilities zones are not affected. For the neutron-rich matter, the δ -meson reduces the size of the spinodal zones for all the magnetic fields considered, contrary to what happens with NL δ .

For $B^* = 5 \times 10^5$ the protons are totally spin polarized in all the models under study and for the complete instability region. The size of the spinodal zone is the largest one.

The existing differences between the different models for neutron rich matter will affect the transition density from a non-homogeneous phase to a homogeneous phase in stellar matter. In fact, the density at the crossing of




FIG. 2: (Color online) Spinodal section in terms of ρ_p versus ρ_n for NL ρ (thin lines) and NL δ (thick lines) at $T = 0$ MeV and for several values of magnetic fields (a) without and (b) with AMM.




FIG. 3: (Color online) Spinodal section in terms of ρ_p versus ρ_n for TW (thin lines) and DDRH δ (thick lines) models at $T = 0$ MeV and for several values of magnetic fields (a) without and (b) with AMM.

the EoS for β -equilibrium stellar matter with the thermodynamic spinodal gives a reasonable prediction of the transition density [25, 26] of the crust to an homogeneous phase in stellar matter. It was shown in [9] that the transition pressure and, at a second level, the transition density define the fraction of the star's moment of inertia contained in the solid crust. In the sequel we will estimate for the different models and magnetic fields the transition density and transition pressure at the inner edge of a compact star in β -equilibrium. We will discuss only the cases for which the crossing is occurring at the first LL. For fields $B^* < 3 \times 10^4$ at densities above the crossing with the first LL, there may occur other crossings with other LL. For these cases the determination of the extension of the non-homogeneous phase should be carried within a more precise method.

The values of the transition density ρ_b^{cross} , and respective proton fraction Y_p and pressure P_m are given for stellar matter under different magnetic field intensities

in Tables II (without AMM) and III (with AMM). For convenience we have plotted the results without AMM in Fig. 4.

For $B = 0$, we can see that the inclusion of the δ -meson reduces the transition density, and the corresponding proton fraction and pressure for NLW models, while for DDRH models the transition density increases and the corresponding proton fraction and pressure decrease. For DDRH models the crossing is occurring at three times larger proton fractions and twice the pressure.

For a finite magnetic field it was shown in [12] for the models TW and TM1 (a parametrization similar to NL ρ) that the transition density increases with the magnetic field. We confirm this result when the δ meson is included and summarize the conclusions:

a) the proton fraction at the crossing increases monotonically with the magnetic field intensity, and for $B^* = 5 \times 10^5$ it reaches the values 0.36 and 0.4 respectively for NLW and DDRH models. The proton fraction does not




FIG. 4: (Color online) Physical quantities at the inner edge of a compact star in the presence of strong magnetic fields: a) density, b) proton fraction and c) pressure.

depend on the inclusion of the δ meson and NLW models predict systematically a smaller proton fraction; b) the transition density is also increasing with the increase for $B^* \leq 5 \times 10^4$. For smaller fields a decrease on the transition density may occur as it is seen with the DDRH models. The smaller the field, the larger the number of bands, defined by the filling of a new LL, will be crossed by the β -equilibrium EOS. As referred before, this situation requires a more carefully study. For the smaller fields the transition densities for NL ρ and NL δ are almost coincident and only above $B^* = 2 \times 10^5$ does the NL δ predict a larger transition density. Looking at the Fig. 1 it is seen that the symmetry energy for these two models start to differ at the saturation density and this may explain the transition density behaviour. However the transition pressure increases faster for NL δ , and for $B^* = 5 \times 10^4$ NL ρ and NL δ have the same transition pressure. For DDRH models, DDRH δ predicts smaller densities for $B^* < 3.5 \times 10^5$. The change of behaviour for the larger fields seems to be related with the change of curvature of the symmetry energy at ~ 0.12 fm $^{-3}$, above which the symmetry energy for DDRH δ becomes much

stiffer. It is interesting to notice that for these models only above $B^* = 5 \times 10^4$ do the transition pressure and density increase with respect to the $B=0$ values; c) even though the proton fractions and transition densities may be similar, all models predict different pressures at the transition. NLW models have larger pressures for $B^* < 3 \times 10^5$, NL δ being the one with the larger values. For the DDRH models and below $B^* < 3 \times 10^5$, it is the model including δ which has the smallest pressure, about half the value calculated with NL δ .

For $B^* = 5 \times 10^5$, the transition density is above the saturation density, $\rho \sim 0.16 - 0.17$ fm $^{-3}$ for NLW models and to $\rho \sim 0.19 - 0.23$ fm $^{-3}$ for DDRH models. For this high field both models with the δ meson predict larger transition densities, pressures and proton fractions due to the stiffness of the symmetry energy at those densities.

TABLE III: Same as Table II, but with AMM included.

B^*	Model	ρ_b^{cross} (fm $^{-3}$)	Y_p	P_m (MeV fm $^{-3}$)
5×10^4	NL ρ	0.079	0.079	0.570
	NL δ	0.079	0.075	0.558
	TW	0.078	0.097	0.606
	DDRH δ	0.080	0.098	0.498
10^5	NL ρ	0.093	0.152	1.274
	NL δ	0.094	0.149	1.329
	TW	0.091	0.178	1.281
	DDRH δ	0.088	0.183	1.108
2×10^5	NL ρ	0.091	0.153	1.206
	NL δ	0.108	0.260	1.819
	TW	0.092	0.305	1.355
	DDRH δ	0.088	0.313	1.198
5×10^5	NL ρ	0.134	0.434	1.693
	NL δ	0.143	0.442	2.245
	TW	0.149	0.488	2.772
	DDRH δ	0.170	0.504	4.356

In Table III we show the same data given in Table II but including the AMM in the calculation. The conclusions are similar: for NLW models, the transition density and the corresponding pressure increase when the δ -meson is included, whereas for DDRH models the inclusion of the isovector-scalar meson decreases the transition density and the corresponding pressure for $B^* < 5 \times 10^5$. In summary, the δ meson gives rise to a larger crust for NLW models and, except to very large fields, to a smaller crust with DDRH models.

The eigenvector associated with the eigenvalue of the free energy curvature matrix defines the direction of the instability and tells us how does the system separate into a dense liquid and a gas phase. It was shown in [13, 27] that in the absence of the magnetic field the direction of instability favors the reduction of the isospin asymmetry of the dense clusters of the system, and increases the isospin asymmetry of the gas surrounding the clusters, the so called distillation effect. This effect is represented

in Fig. 5 where it is seen that for the $B = 0$ curve the fraction $\delta\rho_p^-/\delta\rho_n^-$ is larger than ρ_p/ρ_n below $y_p = 0.5$ and the other way round above. In this figure we show, respectively for NLW models (left) and DDRH models (right), the fraction $\delta\rho_p^-/\delta\rho_n^-$ as a function of y_p for a fixed baryonic density, $\rho = 0.06 \text{ fm}^{-3}$, chosen inside the instability region.

For NLW models and for the two largest fields considered the spinodal region contains a single Landau level and the curve varies smoothly starting at $\delta\rho_p^-/\delta\rho_n^- \sim 1.5$ for NL ρ and ~ 1.62 for NL δ . We point out the very large value of this fraction, always above 1. The magnetic field favors a strong increase of the proton fraction. For $y_p > 0.5$, NL ρ and NL δ behave in a similar way, while below this value the main difference is the larger $\delta\rho_p^-/\delta\rho_n^-$ for NL δ corresponding to a stronger distillation effect.

For $B^* = 10^5$ the spinodal has two bands, see Fig. 2, corresponding to the occupation of the first two Landau levels. The transition from one to the other is clearly seen with a large discontinuity of $\delta\rho_p^-/\delta\rho_n^-$ at $y_p \sim 0.7$. Above this y_p value the curve behaves like the previous ones. However for $y_p < 0.7$ the behavior is quite different: the curve decreases from the value at $y_p=0$, which is independent of the magnitude of the magnetic field, to a value much smaller than the corresponding value of the fraction ρ_p/ρ_n . The same behaviour occurs for NL ρ and NL δ . The fluctuations will not drive the system out of the first Landau level and therefore the larger the proton fraction, the closer the system comes to the top of the band and the smaller are the allowed proton fluctuations. For $y_p > 0.7$ or for the larger magnetic fields the Landau levels are only partially filled and the fluctuations will never drive the system out of the corresponding Landau level. In summary, the effect of the δ -meson on the instability region of NLW models is to reduce the strength of the distillation effect of neutron-rich matter.

For DDRH models the inclusion of the isovector scalar meson reduces the strength of the distillation effect, which is not so dramatic in these models. Including the AMM similar conclusions are drawn. The AMM favors still larger proton fluctuations because neutron polarization stiffens the EOS.

In conclusion, we have studied the influence of the isovector scalar meson on the instabilities of stellar matter under very strong magnetic fields. The fields considered are much stronger than the strongest field measured at the surface of a magnetar which is $B^* \sim 10^2$ for SGR 1806-20 [5]. However, the magnetic fields in the interior of neutrons stars could be larger and the present work shows how fields of the order of $B^* = 10^{18} - 10^{19}$ could affect the inner crust of a compact star. According to the scalar virial theorem [28] the maximum magnetic energy could be comparable to the gravitational energy in an equilibrium configuration, which would correspond to a value of the order of $\sim 10^{18} \text{ G}$.

We have considered four relativistic nuclear models: two models with constant couplings (NL ρ and NL δ) and




FIG. 5: (Color online) $\delta\rho_p^-/\delta\rho_n^-$ plotted as a function of the proton fraction with $\rho = 0.06 \text{ fm}^{-3}$ for the NLW models (NL ρ (thin lines) and NL δ (thick lines)) (left) and DDRH models (TW (thin lines) and DDRH δ (thick lines)) (right) and for several values of the magnetic fields without AMM. The fraction ρ_p/ρ_n is given by the thin dotted line.

two models with density dependent couplings (TW and DDRH δ). Two of the models include the δ -meson. For all the models, we have determined the spinodal surface from the curvature matrix of the free energy for different magnitudes of the magnetic field. It had already been shown [12] that the instability region could be divided in to several bands according to magnitude of the magnetic field and the number of the Landau levels occupied and that the presence of the magnetic field would generally increase the instability region.

We have seen that the inclusion of the δ meson increases (reduces) the size of the thermodynamic instability zone for very neutron-rich matter for the NLW (DDRH) models when compared with the spinodal obtained without the δ -meson. These results reflect themselves on the extension of the crust of a compact star under a strong magnetic field.

By making a rough estimation of the transition density at the inner crust of a compact star under a strong magnetic field from the crossing of the EOS with the thermodynamic spinodal, we have shown that the transition density and associated pressure increases, for NLW models, with the inclusion of the δ meson. On the other hand, for DDRH models the influence of the δ -meson is to decrease the transition density and associated pressure for fields $B^* < 5 \times 10^5$. DDRH δ is predicting the smallest transition densities, almost half the value obtained for NL δ .

If we consider fields $\sim 10^{18} \text{ G}$ ($B^* \sim 5 \times 10^4$), as indicated by the scalar virial theorem, we may take the following conclusions: a) for conventional pulsars TW would predict a larger transition pressure and, therefore, larger fractional moment of inertia in the crust than all the other models, about twice as large; b) for $B^* \sim 5 \times 10^4$, and taking into account AMM, TW would

predict just a small increase of the fractional moment of inertia in the crust, $\sim 15\%$, the NLW models predict an increase of 120% and 220%, respectively, with and without δ -meson, while DDRH δ would predict an increase of 80%. If the AMM are not considered, within TW the fractional moment of inertia in the crust would be smaller than for conventional pulsars, while for DDRH δ there would be a slight increase of 20% and for the NLW models an increase of $\sim 100\%$. The different behaviours in magnetars and conventional pulsars predicted by the different models might be a possibility to impose stronger constraints on the EOS of nuclear matter.

We have also investigated the direction of instability. If the first Landau level is only partially occupied the density fluctuations are such that the system evolves for a state with dense clusters very proton rich immersed in a proton poor gas. A larger proton fraction is favored

energetically due to the degeneracy of the Landau levels. The δ meson will only reduce slightly the strength of this distillation effect for DDRH models and increase it for NLW models. For particles occupying an almost complete Landau level, proton fluctuations are smaller or forbidden and an anti-distillation effect results with a decrease of the proton fraction of the dense clusters.

Acknowledgments

This work was partially supported by FEDER and FCT (Portugal) under Grant SFRH/BPD/14831/2003 and Projects PTDC/FP/64707/2006 and CERN/FP/83505/2008.

-
- [1] V. V. Usov, *Nature* 357, 472 (1992).
 - [2] B. Paczyński, *Acta Astron.* 42, 145 (1992).
 - [3] Christopher Thompson and Robert C. Duncan, *Astrophys. J.* L9, 392 (1992)
 - [4] A. K. Harding and D. Lai, *Rep. Prog. Phys.* 69, 2631 (2006).
 - [5] SGR/APX online Catalogue, <http://www.physics.mcgill.ca/~pulsar/magnetar/main.html>
 - [6] C. Kouveliotou, S. Dieter, T. Strohmayer, J. van Paradijs, G.J. Fishman, C.A. Meegan, K. Hurley, *Nature* 393, 235 (1998).
 - [7] Christopher Thompson and Robert C. Duncan, *Astrophys. J.* 408, 194 (1993)
 - [8] D. G. Ravenhall and C. J. Pethick, *Astrophys. J.* 424, 846 (1994).
 - [9] Bennett Link, Richard I. Epstein, and James M. Lattimer, *Phys. Rev. Lett.* **83**, 3362 (1999).
 - [10] Victoria M. Kaspi, *Astrophys. Space Sci.* 308, 1 (2007).
 - [11] Rim Dib, Victoria M. Kaspi, and Fotis Gavriil, *Astrophys. J.* 673, 1044 (2008).
 - [12] A. Rabhi, C Providência, and J. da Providência, *Phys. Rev C* 79, 015804 (2009).
 - [13] Camille Ducoin, Constança Providência, Alexandre M. Santos, Lucilia Brito, Philippe Chomaz, *Phys. Rev. C* 78, 055801 (2008).
 - [14] S. Kubis and M. Kutschera, *Phys. Lett. B* 399, 191 (1997)
 - [15] B. Liu, V. Greco, V. Baran, M. Colonna, and M. Di Toro, *Phys. Rev. C* 65, 045201 (2002).
 - [16] D. P. Menezes and C. Providência, *Phys. Rev. C* 70, 058801 (2004).
 - [17] S. Typel and H. H. Wolter, *Nucl. Phys. A* 656, 331 (1999).
 - [18] T. Gaitanos, M. Di Toro, S. Typel, V. Baran, C. Fuchs, V. Greco, and H. H. Wolter, *Nucl. Phys. A* 732, 24 (2004).
 - [19] A. Rabhi, C Providência, and J. da Providência, *J. Phys. G: Nucl. Part. Phys.* 35, 125201 (2008).
 - [20] C. Fuchs, H. Lenske, and H. H. Wolter, *Phys. Rev. C* 52, 3043 (1995).
 - [21] S. S. Avancini, L. Brito, D. P. Menezes, and C. Providência, *Phys. Rev. C* 70, 015203 (2004).
 - [22] H. Müller and B. D. Serot, *Phys. Rev. C* 52, 2072 (1995)
 - [23] V. Baran, M. Colonna, M. Di Toro, and A. B. Larionov, *Nucl. Phys. A* 632, 287 (1998).
 - [24] J. Margueron and P. Chomaz, *Phys. Rev. C* **67**, 041602(R) (2003).
 - [25] S.S. Avancini, L. P. Brito, J.R. Marinelli, D.P. Menezes, M.M.W. Moraes, C. Providência and A. M. Santos, *Phys. Rev. C* **79**, 035804 (2009).
 - [26] S.S. Avancini, D.P. Menezes, M.D. Alloy, J.R. Marinelli, M.M.W. Moraes and C. Providência, *Phys. Rev. C* **78**, 015802 (2008).
 - [27] J. Xu, L.W. Chen, B.A. Li and H.R. Ma, *arXiv:0807.4477v1 [nucl-th]*.
 - [28] S. S. Avancini, L. Brito, Ph. Chomaz, D. P. Menezes, and C. Providência, *Phys. Rev. C* 74, 024317 (2006).
 - [29] Dong Lai and Stuart L. Shapiro, *Astrophys. J.* 383, 745 (1991)